arXiv:0804.0701v4 [math.DG] 25 Oct 2008

A_k SINGLARITIES OF WAVE FRONTS.

KENTARO SAJI, MASAAKI UMEHARA, AND KOTARO YAMADA

Dedicated to Professor Yoshiaki Maeda on the occasion of his sixtieth birthday.

Abstract. In this paper, we discuss the recognition problem for A_k-type singularities on wave fronts. We give computable and simple criteria of these singularities, which will play a fundamental role in generalizing the authors' previous work "the geometry of fronts" for surfaces. The crucial point to prove our criteria for A_k-singularities is to introduce a suitable parametrization of the singularities called the "k-th KRSUY-coordinates" (see Section 3). Using them, we can directly construct a versal unfolding for a given singularity. As an application, we prove that a given nondegenerate singular point p on a real (resp. complex) hypersurface (as a wave front) in $\boldsymbol{R}^{n+1}$ (resp. $\boldsymbol{C}^{n+1}$) is differentiably (resp. holomorphically) right-left equivalent to the A_{k+1}-type singular point if and only if the linear projection of the singular set around p into a generic hyperplane $\boldsymbol{R}^n$ (resp. $\boldsymbol{C}^n$) is right-left equivalent to the A_k-type singular point in $\boldsymbol{R}^n$ (resp. $\boldsymbol{C}^n$). Moreover, we show that the restriction of a C^∞-map $f : \boldsymbol{R}^n \to \boldsymbol{R}^n$ to its Morin singular set gives a wave front consisting of only A_k-type singularities. Furthermore, we shall give a relationship between the normal curvature map and the zig-zag numbers (the Maslov indices) of wave fronts.

1. Introduction

Throughout this paper, we denote by $\boldsymbol{K}$ the real number field $\boldsymbol{R}$ or the complex number field $\boldsymbol{C}$. Let m and n be positive integers. A map $f\colon \boldsymbol{K}^m \to \boldsymbol{K}^n$ is called $\boldsymbol{K}$-*differentiable* if it is a C^∞-map when $\boldsymbol{K} = \boldsymbol{R}$, and is a holomorphic map when $\boldsymbol{K} = \boldsymbol{C}$. Let $\langle\ ,\ \rangle_{\boldsymbol{K}}$ be the $\boldsymbol{K}$-*inner product* given by

$$\langle X, Y\rangle_{\boldsymbol{K}} = \sum_{j=1}^{n} x_j y_j \qquad \big(X = (x_1, \ldots, x_n),\ Y = (y_1, \ldots, y_n) \in \boldsymbol{K}^n\big).$$

Let $P^n(\boldsymbol{K})$ be the $\boldsymbol{K}$-projective space and $\boldsymbol{K}^{n+1} \setminus \{\boldsymbol{0}\} \ni X \mapsto [X] \in P^n(\boldsymbol{K})$ the canonical projection. By using the above $\boldsymbol{K}$-inner product, the $\boldsymbol{K}$-projective cotangent bundle $PT^*\boldsymbol{K}^{n+1}$ has the following identification

$$PT^*\boldsymbol{K}^{n+1} = \boldsymbol{K}^{n+1} \times P^n(\boldsymbol{K}),$$

which has the canonical $\boldsymbol{K}$-contact structure. Let $U \subset \boldsymbol{K}^n$ be a domain and

$$L := (f, [\nu]) : U \longrightarrow \boldsymbol{K}^{n+1} \times P^n(\boldsymbol{K})$$

a Legendrian immersion, where ν is a locally defined $\boldsymbol{K}$-differentiable map into $\boldsymbol{K}^{n+1} \setminus \{\boldsymbol{0}\}$ such that

$$\langle df(\boldsymbol{u}), \nu\rangle_{\boldsymbol{K}} = 0 \qquad (\boldsymbol{u} \in TU).$$

Date: October 25, 2008.

2000 *Mathematics Subject Classification.* Primary 57R45; Secondary 57R35, 53D12.

Kentaro Saji was supported by the Grant-in-Aid for Young Scientists (Start-up) No. 19840001, from the Japan Society for the Promotion of Science. Masaaki Umehara and Kotaro Yamada were supported by the Grant-in-Aid for Scientific Research (A) No. 19204005 and Scientific Research (B) No. 14340024, respectively, from the Japan Society for the Promotion of Science.

In this situation, f is called a *wave front* or *front*, and ν is called the $\boldsymbol{K}$*-normal vector field* of f. If $\boldsymbol{K} = \boldsymbol{C}$, $\langle \nu, \nu \rangle_{\boldsymbol{K}}$ might vanish. A point $p \in U$ is called a *singular point* if the front f is not an immersion at p.

In this paper, we shall discuss the recognition problem for A_{k+1}-type singularities on wave fronts. These are fundamental singularities on wave fronts (see [2]). We give a simple and computable necessary and sufficient condition that a given singular point p on a hypersurface (as a wave front) in $\boldsymbol{K}^{n+1}$ is $\boldsymbol{K}$-right-left equivalent to the A_{k+1}-type singular point ($k \leq n$; Theorem 2.4, Corollaries 2.5, 2.6 and 2.8 in Section 2), where two $\boldsymbol{K}$-differentiable map germs $f_j \colon (U_j, p_j) \to (\boldsymbol{K}^{n+1}, p_j')$ $(j = 1, 2)$ are $\boldsymbol{K}$*-right-left equivalent* if there exist $\boldsymbol{K}$-diffeomorphism germs $\psi \colon (U_1, p_1) \to (U_2, p_2)$ and $\Psi : (\boldsymbol{K}^{n+1}, p_1') \to (\boldsymbol{K}^{n+1}, p_2')$ such that $\Psi \circ f_1 = f_2 \circ \psi$ holds. Here, the A_{k+1}-type singularity (or A_{k+1}-front singularity) is a map germ defined by

$$X \longmapsto \left((k+1)t^{k+2} + \sum_{j=2}^{k} (j-1)t^j x_j, -(k+2)t^{k+1} - \sum_{j=2}^{k} j t^{j-1} x_j, X_1 \right) \tag{1.1}$$

at the origin, where $X = (t, x_2, \ldots, x_n)$, $X_1 = (x_2, \ldots, x_n)$. The image of it coincides with the discriminant set $\{F = F_t = 0\} \subset (\boldsymbol{K}^{n+1}; u_0, \ldots, u_n)$ of the versal unfolding

$$F(t, u_0, \ldots, u_n) := t^{k+2} + u_k t^k + \cdots + u_1 t + u_0. \tag{1.2}$$

By definition, A_1-front singularities are regular points. A 3/2-cusp in a plane is an A_2-front singularity and a swallowtail in $\boldsymbol{R}^3$ is an A_3-front singularity.

When $n = 2$, useful criteria for cuspidal edges and swallowtails are given in [7]. We shall give a generalization of the criteria here. The crucial point to prove our criteria for A_{k+1}-singularities is to introduce the "k-th KRSUY-coordinates" as a generalization of the coordinates for cuspidal edges and swallowtails in [7]. Using them, we can directly construct a versal unfolding whose discriminant set coincides with the given singularity.

As an application, when $\boldsymbol{K} = \boldsymbol{R}$, we show that the restriction of a C^∞-map $f : \boldsymbol{R}^{n+1} \to \boldsymbol{R}^{n+1}$ into its Morin singular set (see the appendix) gives a wave front consisting of only A_{k+1}-type singularities ($k \leq n$). Moreover, in the final section, we shall give a relationship between the normal curvature map and the zig-zag numbers (the Maslov indices) of wave fronts.

2. Criteria for A_{k+1}-front singularities

Let $U \subset (\boldsymbol{K}^n; x_1 \ldots, x_n)$ be a domain and $f : U \to \boldsymbol{K}^{n+1}$ a front. Since we will work on a local theory, we may assume that the $\boldsymbol{K}$-normal vector field ν of f is defined on U. We define a $\boldsymbol{K}$-differentiable function λ on U as the determinant

$$\lambda := \det(f_{x_1}, \ldots, f_{x_n}, \nu), \tag{2.1}$$

where $f_{x_j} := \partial f / \partial x_j$ $(j = 1, 2, \ldots, n)$. A point $p \in U$ is 1-*singular* (or *singular*) if $\lambda(p) = 0$, that is, p is a singular point. A point p is 1-*nondegenerate* (or *nondegenerate*) if p is 1-singular and the exterior derivative $d\lambda$ does not vanish at p. The following assertion is obvious:

Lemma 2.1. *The definition of* 1*-nondegeneracy is independent of the choice of local coordinate system* $(x_1, \ldots, x_n)$ *and the choice of a* $\boldsymbol{K}$*-normal vector field* ν*. Moreover, if* p *is* 1*-nondegenerate, the* $\boldsymbol{K}$*-linear map* $df_p : T_pU \to \boldsymbol{K}^{n+1}$ *has a kernel of* $\boldsymbol{K}$*-dimension exactly one.*

If p is 1-nondegenerate, the singular set of $f (=: f_0)$

$$S_1 := S(f) = \{q \in U \,;\, \lambda(q) = 0\}$$

is an embedded $\boldsymbol{K}$-*differentiable* hypersurface of U near p. We denote by TS_1 the $\boldsymbol{K}$-differentiable tangent bundle of $\boldsymbol{K}$-differentiable manifold S_1. Then

$$f_1 := f|_{S_1} : S_1 \longrightarrow \boldsymbol{K}^{n+1}$$

is a $\boldsymbol{K}$-differentiable map. Since $\dim_{\boldsymbol{K}} \ker(df_p) = 1$, we can take a sufficiently small neighborhood V ($\subset U$) of p and non-zero $\boldsymbol{K}$-differentiable vector field η on $S_1 \cap V$ which belongs to the kernel of df, that is $df_q(\eta_q) = 0$ for $q \in S_1 \cap V$. We call η a *null vector field* of f. Moreover, we can construct a $\boldsymbol{K}$-differentiable vector field $\tilde{\eta}$ on V (called an *extended null vector field*) whose restriction $\tilde{\eta}|_{S_1 \cap V}$ on the singular set $S_1 \cap V$ gives a null vector field.

Remark 2.2. Since $\eta \in \ker(df)$ on S_1, we can write the components of η explicitly using determinants of $(n-1)$-submatrices of the Jacobian matrix of f. Then this explicit expression of η gives a $\boldsymbol{K}$-differentiable vector field on a sufficiently small neighborhood of a singular point. Thus, we get an explicit procedure to construct $\tilde{\eta}$. For example, let f be a front $f = (f^1, f^2, f^3, f^4) \colon \boldsymbol{K}^3 \to \boldsymbol{K}^4$ such that $f^1_{x_1} f^2_{x_2} - f^1_{x_2} f^2_{x_1} \neq 0$. Then

$$\tilde{\eta} = \left(\det \begin{pmatrix} f^1_{x_2} & f^1_{x_3} \\ f^2_{x_2} & f^2_{x_3} \end{pmatrix}, -\det \begin{pmatrix} f^1_{x_1} & f^1_{x_3} \\ f^2_{x_1} & f^2_{x_3} \end{pmatrix}, \det \begin{pmatrix} f^1_{x_1} & f^1_{x_2} \\ f^2_{x_1} & f^2_{x_2} \end{pmatrix} \right)$$

is an extended null vector field.

Let $\lambda' \colon U \to \boldsymbol{K}$ and $f', f'' : U \to \boldsymbol{K}^{n+1}$ be the derivatives $d\lambda(\tilde{\eta})$, $df(\tilde{\eta})$ and $df'(\tilde{\eta})$, respectively. Then by definition, we have

$$S_1 := \{q \in U \,;\, f'(q) = \mathbf{0}\}, \tag{2.2}$$

where $\mathbf{0} = (0, \ldots, 0)$. The following assertion holds:

Lemma 2.3. *The singular set* $S_2 := S(f_1)$ *of* $f_1 \colon S_1 \to \boldsymbol{K}^{n+1}$ *has the following expressions*:

$$\begin{aligned} S_2 &= \{q \in S_1 \,;\, \eta_q \in T_q S_1\} = \{q \in S_1 \,;\, \lambda'(q) = 0\} = \{q \in U \,;\, \lambda(q) = \lambda'(q) = 0\} \\ &= \{q \in S_1 \,;\, f''(q) = \mathbf{0}\} = \{q \in U \,;\, f'(q) = f''(q) = \mathbf{0}\}. \end{aligned}$$

Proof. A point $q \in S_1$ belongs to S_2 if and only if there exists $\xi \in T_q S_1 \setminus \{\mathbf{0}\}$ such that $df_1(\xi) = 0$. Since $\dim_{\boldsymbol{K}} \ker(df_q) = 1$, the vector ξ must be proportional to η_q. Thus we get $S_2 = \{q \in S_1 \,;\, \eta_q \in T_q S_1\}$. Since S_1 is a level set of λ, $\eta_q \in T_q S_1$ if and only if $d\lambda(\eta_q) = \lambda'(q) = 0$, which proves the first equation. By (2.2) and by $\eta \in TS_1$ on S_2, the derivative $f'' = df'(\eta)$ of f' with respect to η vanishes on S_2, that is $S_2 \subset \{q \in S_1 \,;\, f''(q) = \mathbf{0}\}$. Next we suppose $f''(q) = \mathbf{0}$ ($q \in S_1$). Since $\lambda' = 0$ on S_2, by taking a new coordinate system $(z_1, \ldots, z_n)$ on U such that $\partial/\partial z_n = \tilde{\eta}$, we have

$$\lambda'(q) = \{\varphi \det(f_{z_1}, \ldots, f_{z_{n-1}}, f', \nu)\}'(q) = \varphi(q) \det(f_{z_1}, \ldots, f_{z_{n-1}}, f'', \nu)(q) = 0,$$

where $\varphi := \det(\partial z_i / \partial x_j)$ is the Jacobian. Consequently $q \in S_2$ holds, that is $S_2 = \{q \in S_1 \,;\, f''(q) = \mathbf{0}\}$. □

A 1-nondegenerate singular point $p \in S_1$ is called 2-*singular* if $p \in S_2$. Moreover, a 2-singular point $p \in S_2$ is 2-*nondegenerate* if $(d\lambda')_p \neq 0$ on $T_p S_1$. In this case, S_2 is an embedded hypersurface of S_1 around p if it is 2-nondegenerate.

Now, we define the j-singularity and the j-nondegeneracy inductively ($2 \leq j \leq n$): Firstly, we give the following notations:

$$\begin{aligned} &\lambda^{(0)} := \lambda, \qquad \lambda^{(1)} := \lambda', \qquad \lambda^{(l)} := d\lambda^{(l-1)}(\tilde{\eta}), \\ &f^{(0)} := f, \qquad f^{(1)} := f', \qquad f^{(l)} := df^{(l-1)}(\tilde{\eta}), \end{aligned}$$

where $l = 1, 2, 3, \ldots$. We fix a $(j-1)$-nondegenerate singular point p of f. Then the $(j-1)$-st singular set

$$S_{j-1} := S(f_{j-2})$$

is an embedded hypersurface of S_{j-2} around p. (Here we replace U by a sufficiently small neighborhood of p at each induction step if necessary. In particular, we may assume $S_0 := U$.) We set

$$f_{j-1} := f|_{S_{j-1}} : S_{j-1} \longrightarrow \boldsymbol{K}^{n+1}.$$

As in the proof of Lemma 2.3, by an inductive use of the identity

$$\begin{aligned}\lambda^{(j-1)}(q) &= \{\varphi \det(f_{z_1}, \ldots, f_{z_{n-1}}, f', \nu)\}^{(j-1)}(q)\\ &= \varphi(q)\det(f_{z_1}, \ldots, f_{z_{n-1}}, f^{(j)}, \nu)(q),\end{aligned}$$

$S_j := S(f_{j-1})$ satisfies

$$\begin{aligned}S_j &= \{q \in S_{j-1}\,;\, \eta_q \in T_q S_{j-1}\}\\ &= \{q \in S_{j-1}\,;\, \lambda^{(j-1)}(q) = 0\} = \{q \in U\,;\, \lambda(q) = \cdots = \lambda^{(j-1)}(q) = 0\}\\ &= \{q \in S_{j-1}\,;\, f^{(j)}(q) = \boldsymbol{0}\} = \{q \in U\,;\, f'(q) = \cdots = f^{(j)}(q) = \boldsymbol{0}\}.\end{aligned} \tag{2.3}$$

Each point of S_j is called a *j-singular point*. A point $q \in S_j$ is called *j-nondegenerate* if $(d\lambda^{(j-1)})_q \neq 0$ on $T_q S_{j-1}$, that is

$$T_q S_{j-1} \not\subset \ker(d\lambda^{(j-1)})_q. \tag{2.4}$$

These definitions do not depend on the choice of a coordinate system $(x_1, \ldots, x_n)$ of U. Under these notations, a criterion for A_{k+1}-front singularities is stated as follows:

Theorem 2.4. *Let $f : (U, p) \to (\boldsymbol{K}^{n+1}, f(p))$ be a front, where U is a domain in $\boldsymbol{K}^n$. Then f at p is $\boldsymbol{K}$-right-left equivalent to the A_{k+1}-front singularity if and only if p is k-nondegenerate but is not $(k+1)$-singular $(k \leq n)$.*

Though k-singular points are defined only for $k \leq n$, we define any points are not $(n+1)$-singular. This theorem is proved in Sections 3 and 4.

Corollary 2.5. *Let $f : U \to \boldsymbol{K}^{n+1}$ be a front. Then f at $p \in U$ is $\boldsymbol{K}$-right-left equivalent to the A_{k+1}-front singularity $(1 \leq k \leq n)$ if and only if*

$$\lambda = \lambda' = \cdots = \lambda^{(k-1)} = 0, \quad \lambda^{(k)} \neq 0 \tag{2.5}$$

at p and the Jacobian matrix of $\boldsymbol{K}$-differentiable map $\Lambda := (\lambda, \lambda', \ldots, \lambda^{(k-1)}) : U \to \boldsymbol{K}^k$ is of rank k at p.

Corollary 2.6. *Let U be a domain in $\boldsymbol{K}^n$, $f\colon U \to \boldsymbol{K}^{n+1}$ a front, and $p \in U$ a nondegenerate singular point. Take a local tangent frame field $\{v_1, \ldots, v_{n-1}\}$ of the singular set $S(f)$ (a smooth hypersurface in U) around p and set*

$$\mu := \det(v_1, \ldots, v_{n-1}, \eta), \tag{2.6}$$

as the determinant function on $\boldsymbol{K}^n$ where η is a null vector field. Then p is 2-singular if and only if $\mu(p) = 0$. In particular, p is $\boldsymbol{K}$-right-left equivalent to the A_2-singularity if and only if $\mu(p) \neq 0$. Moreover, a 2-singular point p is 2-nondegenerate if and only if $d\mu(T_p S(f)) \neq \{0\}$. Furthermore, p is $\boldsymbol{K}$-right-left equivalent to the A_3-singularity if and only if $\mu(p) = 0$ and $d\mu(\eta_p) \neq 0$ hold. (If $\mu(p) = 0$, then the null vector η_p at p is tangent to $S(f)$ and $d\mu(\eta_p)$ is well-defined.)

Remark 2.7. These assertions for $n = 2$ have been already proved in [7]. The criterion for A_2-singularities for general n has been shown in [10].

Let $f\colon U \to \boldsymbol{K}^{n+1}$ a front, and $p \in U$ a 2-nondegenerate singular point. Then $S_2 = S(f_1)$ is a submanifold of codimension 2 in U and Lemma 2.3 yields that the null vector field η is a tangent vector field of S_2, and we can set

$$\mu' = d\mu(\eta), \quad \mu'' = d\mu'(\eta), \quad \cdots, \quad \mu^{(k-1)} = d\mu^{(k-2)}(\eta).$$

Then these functions $\mu', \mu'' \cdots, \mu^{(k-1)}$ are C^∞-functions on S_2.

Corollary 2.8. *Let U be a domain in $\boldsymbol{K}^n$, $f\colon U \to \boldsymbol{K}^{n+1}$ a front, and $p \in U$ a 2-nondegenerate singular point. Then p is $\boldsymbol{K}$-right-left equivalent to the A_{k+1}-singularity $(3 \leq k \leq n)$ if and only if*

$$\mu = \mu' = \cdots = \mu^{(k-2)} = 0, \qquad \mu^{(k-1)} \neq 0 \qquad \text{at } p \tag{2.7}$$

hold, and the Jacobian matrix of the map $\Phi_\mu := (\mu', \ldots, \mu^{(k-2)})\colon S_2 \to \boldsymbol{K}^{k-2}$ is of rank $k-2$ at p.

Remark 2.9. In Corollary 2.8, the criterion for A_4-singularities reduces to 2-nondegeneracy and

$$\mu(p) = \mu'(p) = 0, \qquad \mu''(p) \neq 0,$$

since the condition $\mu''(p) \neq 0$ implies that the map $\Phi_\mu = \mu'$ is of rank 1 at p.

Proofs of these assertions are given in Section 4.

Let $f : U \to \boldsymbol{K}^{n+1}$ be a front and p a 1-nondegenerate singular point. Then we can consider the restriction $f_1 : S(f) \to \boldsymbol{K}^{n+1}$ of f into S_1. Let us denote the limiting tangent hyperplane by

$$\nu(p)^\perp := \{\boldsymbol{v} \in T_{f(p)}\boldsymbol{K}^{n+1}\,;\, \langle \boldsymbol{v}, \nu(p)\rangle_{\boldsymbol{K}} = 0\}.$$

The following assertion can be proved straightforwardly.

Corollary 2.10. *Let $f : U \to \boldsymbol{K}^{n+1}$ be a front and $p \in U$ a 1-nondegenerate singular point. For any vector $\boldsymbol{n} \notin \nu(p)^\perp$, the following are equivalent*:

(1) *f at p is an A_{k+1}-front singularity $(k \leq n)$.*
(2) *The projection $\pi_{\boldsymbol{n}} \circ f|_{S(f)}$ at p is an A_k-front singularity $(k \leq n)$.*

Here, $\pi_{\boldsymbol{n}}$ is the normal projection with respect to $\boldsymbol{n}$ to the hyperplane

$$\boldsymbol{n}^\perp := \{\boldsymbol{v} \in T_{f(p)}\boldsymbol{K}^{n+1}\,;\, \langle \boldsymbol{v}, \boldsymbol{n}\rangle_{\boldsymbol{K}} = 0\}$$

and an A_1-front singularity means a regular point.

The following assertion follows immediately from Theorem A.1 in the appendix.

Corollary 2.11. *Let Ω be a domain in $\boldsymbol{K}^{n+1}$ and $f : \Omega \to \boldsymbol{K}^{n+1}$ a $\boldsymbol{K}$-differentiable map. Suppose that $p \in \Omega$ is a 1-nondegenerate singular point, namely, the exterior derivative of the Jacobian of f does not vanish at p. Then the following are equivalent*:

(1) *p is an A_k-Morin singular point of f.*
(2) *$f|_{S(f)}$ is a front, and p is an A_k-front singularity of $f|_{S(f)}$.*

Here, the A_k-Morin singularities are defined in the appendix, and the A_0-Morin singularity means a regular point.

3. Adopted coordinates and k-nondegeneracy

In [7], a certain kind of special coordinate system was introduced to treat the recognition problem of cuspidal edges and swallowtails in $\boldsymbol{R}^3$. In this section, we give a generalization of such a coordinate system around a singular point, which will play a crucial role.

Lemma 3.1 (Existence of the k-th KRSUY-coordinate system)**.** *Let $f : U \to \boldsymbol{K}^{n+1}$ be a front and $p \in U$ a k-nondegenerate singular point ($1 \le k \le n$). Then there exists a $\boldsymbol{K}$-differentiable coordinate system $(z_1, \ldots, z_n)$ around p and a (non-degenerate) $\boldsymbol{K}$-affine transformation $\Theta : \boldsymbol{K}^{n+1} \to \boldsymbol{K}^{n+1}$ such that*

$$\hat{f}(z_1, \ldots, z_n) = \left(\hat{f}^1, \ldots, \hat{f}^{n+1}\right) = \Theta \circ f(z_1, \ldots, z_n)$$

satisfies the following properties:

- (P_0) *The point p corresponds to $(z_1, \ldots, z_n) = (0, \ldots, 0)$.*
- (P_1) *$\partial_{z_n} := \partial/\partial z_n$ gives an extended null vector field on U.*
- (P_2) *If $k \ge 2$, then the $\boldsymbol{K}$-vector space T_pS_i $(i = 1, \ldots, k-1)$ is spanned by $\partial_{z_{i+1}}, \ldots, \partial_{z_n}$, that is*

 $$T_pS_i = \mathrm{Span}_{\boldsymbol{K}}\left\{\partial_{z_{i+1}}, \ldots, \partial_{z_n}\right\},$$

 where $\partial_{z_j} := \partial/\partial z_j$, $j = 1, \ldots, n$.
- (P_3) *Suppose $k < n$. If p is $(k+1)$-singular, then $T_pS_k = \mathrm{Span}_{\boldsymbol{K}}\left\{\partial_{z_{k+1}} \ldots, \partial_{z_n}\right\}$. On the other hand, if p is not $(k+1)$-singular, $T_pS_k = \mathrm{Span}_{\boldsymbol{K}}\left\{\partial_{z_k}, \ldots, \partial_{z_{n-1}}\right\}$.*
- (P_4) *$\hat{f}(p) = \boldsymbol{0}$ and $\nu(p) = (1, 0, \ldots, 0)$.*
- (P_5) *$\hat{f}(z_1, \ldots, z_n) = \left(\hat{f}^1(z_1, \ldots, z_n), \hat{f}^2(z_1, \ldots, z_n), z_1, \ldots, z_{n-1}\right)$ for $n \ge 2$ and $\hat{f}(z_1) = \left(\hat{f}^1(z_1), \hat{f}^2(z_1)\right)$ for $n = 1$.*
- (P_6) *If $n \ge 2$, $(\hat{f}^1)_{z_i}(p) = (\hat{f}^2)_{z_i}(p) = 0$ holds for $i = 1, \ldots, n-1$.*

We call a coordinate system on U satisfying the properties (P_0), (P_1), (P_2) and (P_3) a "*k-adopted coordinate system*", and we call by the "*k-th KRSUY-coordinate system*" a pair of coordinate systems on U and $\boldsymbol{K}^{n+1}$ satisfying all above conditions. (In [7], the existence of the coordinates for $k = 1$, 2 and $n = 2$ was shown.)

Proof of Lemma 3.1. By replacing U to be a smaller neighborhood if necessary, we can take an extended null vector field $\tilde{\eta}$ on U (see Remark 2.2). Now we assume that $p \in S_k$ is k-nondegenerate. Then S_k is a submanifold of U of codimension k. So we can take a basis $\{\xi_1, \ldots, \xi_{n-1}, \tilde{\eta}_p\}$ of T_pU such that

$$\begin{cases} \mathrm{Span}_{\boldsymbol{K}}\{\xi_k, \ldots, \xi_{n-1}\} = T_pS_k & \text{if } k < n \text{ and } p \text{ is not } k\text{-singular}, \\ \mathrm{Span}_{\boldsymbol{K}}\{\xi_{k+1}, \ldots, \xi_{n-1}, \tilde{\eta}_p\} = T_pS_k & \text{if } k < n \text{ and } p \text{ is } k\text{-singular}, \end{cases}$$

and

$$\mathrm{Span}_{\boldsymbol{K}}\{\xi_{i+1}, \ldots, \xi_{n-1}, \tilde{\eta}_p\} = T_pS_i \qquad (1 \le i \le k-1 \quad \text{and} \quad k \ge 2).$$

We now take a local coordinate system $(s_1, s_2, \ldots, s_n)$ around p such that $\partial_{s_i}(p) = \xi_i$ $(i = 1, \ldots, n-1)$, and consider the following two involutive distributions T_1 and T_2 of rank $n-1$ and 1 respectively

$$T_1(x) := \mathrm{Span}_{\boldsymbol{K}}\left\{\partial_{s_1}(x), \partial_{s_2}(x), \ldots, \partial_{s_{n-1}}(x)\right\}, \quad T_2(x) := \mathrm{Span}_{\boldsymbol{K}}\{\tilde{\eta}_x\} \quad (x \in V),$$

where V is a sufficiently small neighborhood of p. Since two distributions $T_1(p)$ and $T_2(p)$ span T_pV, the lemma in [6, page 182] yields the existence of local coordinate system $(w_1, w_2, \ldots, w_n)$ around p such that

$$T_1 := \mathrm{Span}_{\boldsymbol{K}}\left\{\partial_{w_1}, \partial_{w_2}, \ldots, \partial_{w_{n-1}}\right\}, \qquad \partial_{w_n} \in \mathrm{Span}_{\boldsymbol{K}}\{\tilde{\eta}\}\,(= T_2).$$

Moreover, by a suitable affine transformation of $\boldsymbol{K}^n$, we may assume that $w_i(p) = 0$ $(i = 1, \ldots, n)$ and $\partial_{w_i}(p) = \xi_i$ $(i = 1, \ldots, n-1)$. Replacing U to be sufficiently smaller, we may assume that this new coordinate system $(w_1, \ldots, w_n)$ is defined on

U. Moreover, we may reset $\tilde{\eta} = \partial_{w_n}$. Thus $(U; w_1, \ldots, w_n)$ satisfies (P_0), (P_1), (P_2) and (P_3). By a suitable affine transformation of $\boldsymbol{K}^{n+1}$, we may assume that

$$f(p) = \mathbf{0}, \quad \nu(p) = \boldsymbol{e}_1, \qquad \frac{\partial f}{\partial w_i}(p) = \boldsymbol{e}_{i+2} \quad (i = 1, \ldots, n-1), \tag{3.1}$$

where $\boldsymbol{e}_j$ denotes the j-th canonical vector

$$\boldsymbol{e}_j := (0, \ldots, 0, 1, 0, \ldots, 0) \qquad (j = 1, 2, \ldots, n+1) \tag{3.2}$$

of $\boldsymbol{K}^{n+1}$. We set $f = (f^1, \ldots, f^{n+1})$. Then it holds that

$$\frac{\partial f^{j+2}}{\partial w_l}(p) = \delta^j_l \qquad (j, l = 1, \ldots, n-1), \tag{3.3}$$

where δ^j_l is Kronecker's delta. Define new functions by

$$g_j(z_1, \ldots, z_{n-1}; w_1, \ldots, w_n) := f^{j+2}(w_1, \ldots, w_n) - z_j \qquad (j = 1, \ldots, n-1).$$

Since $\partial g_j/\partial w_l(0) = \delta^j_l$ for $(j, l = 1, \ldots, n-1)$, the implicit function theorem yields that there exist functions $\varphi_j(z_1, \ldots, z_{n-1}, w_n)$ $(j = 1, \ldots, n-1)$ such that

$$g_j\big(z_1, \ldots, z_{n-1}, \varphi_1(z_1, \ldots, z_{n-1}, w_n), \ldots, \varphi_{n-1}(z_1, \ldots, z_{n-1}, w_n), w_n\big) = 0.$$

If we set $w_n := z_n$ and

$$w_j := \varphi_j(z_1, \ldots, z_{n-1}, w_n) \qquad (j = 1, \ldots, n-1),$$

then $(z_1, \ldots, z_n)$ gives a new local coordinate system of U around p such that

$$z_j = f^{j+2}\big(\varphi_1(z_1, \ldots, z_{n-1}, z_n), \ldots, \varphi_{n-1}(z_1, \ldots, z_{n-1}, z_n), z_n\big), \tag{3.4}$$

$$\frac{\partial w_j}{\partial z_l}(p) = \delta^j_l \qquad (j = 1, \ldots, n-1,\ l = 1, \ldots, n). \tag{3.5}$$

Since ∂_{w_n} is the null direction, we have $\partial f/\partial w_n = 0$. Differentiating (3.4) with respect to z_n, we have

$$0 = \sum_{l=1}^{n-1} \frac{\partial f^{j+2}}{\partial w_l}\frac{\partial \varphi_l}{\partial z_n} + \frac{\partial f^{j+2}}{\partial w_n} = \sum_{l=1}^{n-1} \frac{\partial f^{j+2}}{\partial w_l}\frac{\partial \varphi_l}{\partial z_n} \qquad \text{on } S_1.$$

Since the matrix $\big(\partial f^{j+2}/\partial w_l\big)_{j,l=1,\ldots,n-1}$ is regular near p by (3.3),

$$\frac{\partial w_l}{\partial z_n} = \frac{\partial \varphi_l}{\partial z_n} = 0 \qquad (\text{on } S_1 \text{ near } p)$$

hold for $l = 1, \ldots, n-1$. In particular, (3.5) holds for $j = n$.

Thus if we set

$$\hat{f}(z_1, \ldots, z_n) := f\big(\varphi_1(z_1, \ldots, z_n), \ldots \varphi_{n-1}(z_1, \ldots, z_n), z_n\big),$$

then

$$\frac{\partial \hat{f}}{\partial z_n} = \sum_{j=1}^{n-1} \frac{\partial f}{\partial w_j}\frac{\partial \varphi_j}{\partial z_n} + \frac{\partial f}{\partial w_n} = \mathbf{0} \qquad \text{on } S_1 \text{ near } p,$$

holds which implies (P_1). By (3.5), $\partial_{z_j}(p) = \partial_{w_j}(p)$ for $j = 1, \ldots, n$. Hence we have (P_2) and (P_3). Moreover,

$$\frac{\partial \hat{f}}{\partial z_i}(p) = \left(\sum_{j=1}^{n-1} \frac{\partial f}{\partial w_j}(p)\frac{\partial w_j}{\partial z_i}(p)\right) + \frac{\partial f}{\partial w_n}(p) = \frac{\partial f}{\partial w_i}(p) = \boldsymbol{e}_{i+2},$$

which implies (P_6). Now (P_4) and (P_5) follow immediately from (3.1) and (3.4) respectively. □

Now we use the notation $' = \partial/\partial z_n$ under the k-adopted coordinate system.

Lemma 3.2. *Let p be a k-nondegenerate singular point of f. Under a k-th adopted coordinate system $(z_1, \ldots, z_n)$, the following hold*:

(1) *If $k = 1$, then p is 2-singular if and only if $\lambda'(p) = 0$. Moreover $p(\in S_2)$ is 2-nondegenerate if and only if*

$$\big(\lambda'_{z_2}(p), \ldots, \lambda'_{z_{n-1}}(p), \lambda''(p)\big) \neq \mathbf{0}.$$

(2) *If $k \geq 2$, then $\lambda^{(m-2)}_{z_{m-1}}(p) \neq 0$ and*

$$\lambda^{(m-2)}_{z_m}(p) = \cdots = \lambda^{(m-2)}_{z_{n-1}}(p) = \lambda^{(m-1)}(p) = 0 \tag{3.6}$$

hold for $m = 2, \ldots, k$. Moreover p is $(k+1)$-singular $(k < n)$ if and only if $\lambda^{(k)}(p) = 0$. Furthermore $p(\in S_{k+1})$ is $(k+1)$-nondegenerate if and only if

$$\big(\lambda^{(k)}_{z_{k+1}}(p), \ldots, \lambda^{(k)}_{z_{n-1}}(p), \lambda^{(k+1)}(p)\big) \neq \mathbf{0}.$$

Proof. When $1 \leq k < n$, p is $(k+1)$-singular if and only if $\lambda^{(k)}(p) = 0$ by (2.3). Suppose now p is $(k+1)$-singular $(k < n)$. By definition, p is $(k+1)$-nondegenerate if and only if $T_pS_k \not\subset \ker(d\lambda^{(k)})_p$. Since $\mathrm{Span}_{\boldsymbol{K}}\{\partial_{z_{k+1}}, \ldots, \partial_{z_n}\} = T_pS_k$ by (P_2), this is equivalent to $\big(\lambda^{(k)}_{z_{k+1}}, \ldots, \lambda^{(k)}_{z_{n-1}}, \lambda^{(k+1)}\big)(p) \neq \mathbf{0}$, which proves (1) and the latter part of (2). Next, we assume $2 \leq m \leq k \leq n$ and prove the first part of (2): By (2.3), we have $S_{m-1} = \{q \in S_{m-2}\,;\, \lambda^{(m-2)}(q) = 0\}$. By the assumption (P_2), we have $T_pS_{m-1} = \mathrm{Span}_{\boldsymbol{K}}\{\partial_{z_m}, \ldots, \partial_{z_n}\}$. In particular, $\lambda^{(m-2)}$ is constant along these directions, which implies (3.6). On the other hand, since $k \geq m \geq 2$, p is $(m-1)$-nondegenerate. By (2.4), we have $T_pS_{m-2} \not\subset \ker(d\lambda^{(m-2)})_p$, which implies $\lambda^{(m-2)}_{z_{m-1}}(p) \neq 0$ because of (3.6). □

Lemma 3.3. *Suppose that $p \in U$ is a singular point of f which is k-nondegenerate but not $(k+1)$-singular. Then under the k-th adopted coordinate system, $f^{(k+1)}(p) \neq \mathbf{0}$ holds. Moreover, if $k \geq 2$, then $f^{(m-1)}_{z_{m-1}}(p) \neq \mathbf{0}$ and*

$$f^{(m-1)}_{z_m}(p) = \cdots = f^{(m-1)}_{z_{n-1}}(p) = f^{(m)}(p) = \mathbf{0} \tag{3.7}$$

hold for $m = 2, \ldots, k$.

Proof. Since $p \in S_k \setminus S_{k+1}$, (2.3) yields that

$$f'(p) = \cdots = f^{(k)}(p) = \mathbf{0}, \quad f^{(k+1)}(p) \neq \mathbf{0}.$$

Suppose that $k \geq m \geq 2$. Then p is m-nondegenerate. By (2.3), the map $f^{(m-1)}$ vanishes along S_{m-1}. Since T_pS_{m-1} is spanned by $\partial_{z_m}, \ldots, \partial_{z_n}$ because of (P_2), we have (3.7). On the other hand, since $f' = \cdots = f^{(m-1)} = \mathbf{0}$ at p, Lemma 3.2 yields that

$$0 \neq \lambda^{(m-2)}_{z_{m-1}}(p) = \det(f_{z_1}, \cdots, f_{z_{n-1}}, f^{(m-1)}_{z_{m-1}}, \nu)(p),$$

which implies that $f^{(m-1)}_{z_{m-1}}(p) \neq \mathbf{0}$. □

4. Proof of the criteria

Let $f\colon U \to \boldsymbol{K}^{n+1}$ be a front and ν be a $\boldsymbol{K}$-normal vector. Let $p \in U$ be a singular point of f which is k-nondegenerate but not $(k+1)$-singular. We denote the k-th KRSUY coordinates by $(U; z_1, \ldots, z_n)$ and $(\boldsymbol{K}^{n+1}; x_1, \ldots, x_{n+1})$. Set

$$X := (x_1, x_2), \quad Y := (z_1, \ldots, z_{n-1}), \quad (X, Y) = (x_1, x_2, z_1, \ldots, z_{n-1}).$$

We define a function $\Phi\colon \boldsymbol{K}^{n+1} \times \boldsymbol{K} \to \boldsymbol{K}$ by

$$\Phi(X, Y, z_n) := \langle \nu(Y, z_n), f(Y, z_n) - (X, Y) \rangle_{\boldsymbol{K}}\,. \tag{4.1}$$

Then we have the following

Proposition 4.1. *The discriminant set*

$$D_\Phi := \left\{ (X,Y) \in \boldsymbol{K}^{n+1} \,;\, \Phi(X,Y,t) = \frac{\partial \Phi}{\partial t}(X,Y,t) = 0 \text{ for some } t \right\}$$

coincides with the image $\mathrm{Im}(f)$ *of* f.

Proof. One can easily prove that $D_\Phi \supset \mathrm{Im}(f)$. Next we show $D_\Phi \subset \mathrm{Im}(f)$. Suppose $(X,Y) \in D_\Phi$. Since $\langle f_{z_n}, \nu \rangle_{\boldsymbol{K}} = 0$, $(X,Y) \in D_\Phi$ is equivalent to

$$\langle \nu, f - (X,Y) \rangle_{\boldsymbol{K}} = 0 \quad \text{and} \quad \langle \nu', f - (X,Y) \rangle_{\boldsymbol{K}} = 0. \tag{4.2}$$

Since we are using the k-th KRSUY-coordinates, (4.2) reduces to

$$\left\langle \nu, (f^1 - x_1, f^2 - x_2, 0, \ldots, 0) \right\rangle_{\boldsymbol{K}} = \left\langle \nu', (f^1 - x_1, f^2 - x_2, 0, \ldots, 0) \right\rangle_{\boldsymbol{K}} = 0. \tag{4.3}$$

By the following Lemma 4.2, the first two components of ν, ν' are linearly independent near p as vectors in $(\boldsymbol{K}^2; x_1, x_2)$. Hence by (4.3), we have $x_1 = f^1$, $x_2 = f^2$. Since

$$(X,Y) = (f^1(Y, z_n), f^2(Y, z_n), Y) = f(Y, z_n) \in \mathrm{Im}(f)$$

holds, we have $D_\Phi \subset \mathrm{Im}(f)$. □

Lemma 4.2. *The vectors* $f_{z_1}, \ldots, f_{z_{n-1}}, \nu, \nu'$ *are linearly independent near* p. *Moreover, if we write* $\nu = (\nu^1, \ldots, \nu^{n+1})$, *then* (ν^1, ν^2) *and* $\big((\nu^1)', (\nu^2)'\big)$ *are linearly independent in* $\boldsymbol{K}^2$ *near* p.

Proof. Clearly, $f_{z_1}, \ldots, f_{z_{n-1}}, \nu$ are linearly independent. Since f is a front and $f'(p) = \boldsymbol{0}$, we have $\nu' \notin \mathrm{Span}_{\boldsymbol{K}}\{\nu\}$. We assume $\nu' \in \mathrm{Span}_{\boldsymbol{K}}\{f_{z_1}, \ldots, f_{z_{n-1}}, \nu\}$ at p. Since $\langle \nu', f_{z_i} \rangle_{\boldsymbol{K}} = -\langle \nu, f'_{z_i} \rangle_{\boldsymbol{K}} = \langle \nu_{z_i}, f' \rangle_{\boldsymbol{K}} = 0$, $\nu' \in \mathrm{Span}_{\boldsymbol{K}}\{\nu\}$, a contradiction. Moreover, since $f_{z_j}(p) = \boldsymbol{e}_{j+2}$ by (P_5), (ν^1, ν^2) and $\big((\nu^1)', (\nu^2)'\big)$ are linearly independent. Then the second part of the lemma follows. □

Lemma 4.3. *Suppose that* $p \in S(f)$ *is* k*-nondegenerate but not* $(k+1)$*-singular. Under the* k*-th KRSUY-coordinates at* p, *it holds that*

1. $\Phi^{(m)}(\boldsymbol{0}, 0) = 0$ $(m = 0, \ldots, k+1)$ *and* $\Phi^{(k+2)}(\boldsymbol{0}, 0) \neq 0$, *where* $\boldsymbol{0}$ *is the origin of* $\boldsymbol{K}^{n+1}$,
2. *the Jacobian matrix of the map* $(\Phi, \Phi', \ldots, \Phi^{(k)}) : \boldsymbol{K}^{n+2} \to \boldsymbol{K}^{k+1}$ *is of rank* $k+1$ *at the origin.*

Proof. By identifying $\bigwedge^n \boldsymbol{K}^{n+1}$ with $\boldsymbol{K}^{n+1}$ with respect to $\langle\ ,\ \rangle_{\boldsymbol{K}}$, $f_{z_1}, \ldots, f_{z_{n-1}}, \nu$ and $f_{z_1} \wedge \cdots \wedge f_{z_{n-1}} \wedge \nu$ are linearly independent near p, where $\wedge$ denotes the exterior product. Then, one can write

$$\nu' = -a f_{z_1} \wedge \cdots \wedge f_{z_{n-1}} \wedge \nu + \left(\sum_{l=1}^{n-1} b_l f_{z_l} \right) + c\nu, \tag{4.4}$$

for some functions a, b_1, …, b_{n-1}, c in $(z_1, \ldots, z_n)$. By Lemma 4.2, we have

$$a \neq 0 \qquad \text{near } p. \tag{4.5}$$

Here, on the singular set S_1, we have

$$0 = \langle \nu_{z_j}, f' \rangle_{\boldsymbol{K}} = \langle \nu', f_{z_j} \rangle_{\boldsymbol{K}} = \sum_{l=1}^{n-1} b_l \langle f_{z_l}, f_{z_j} \rangle_{\boldsymbol{K}} \qquad (j = 1, \ldots, n-1).$$

Hence we have

$$b_j = 0 \quad (j = 1, \ldots, n-1) \qquad \text{on } S_1 \text{ near } p. \tag{4.6}$$

By the definition of λ, we can write

$$\langle \nu', f' \rangle_{\boldsymbol{K}} = a\lambda + \sum_{l=1}^{n-1} b_l \langle f_{z_l}, f' \rangle_{\boldsymbol{K}}. \tag{4.7}$$

Since $\langle \nu', f' \rangle_{\boldsymbol{K}} = 0$, differentiating (4.1) with respect to z_n, we have

$$\begin{aligned} \Phi' &= \langle \nu', f - (X, Y) \rangle_{\boldsymbol{K}} + \langle \nu, f' \rangle_{\boldsymbol{K}} = \langle \nu', f - (X, Y) \rangle_{\boldsymbol{K}} , \\ \Phi'' &= \langle \nu'', f - (X, Y) \rangle_{\boldsymbol{K}} + 2 \langle \nu', f' \rangle_{\boldsymbol{K}} + \langle \nu, f'' \rangle_{\boldsymbol{K}} \\ &= \langle \nu'', f - (X, Y) \rangle_{\boldsymbol{K}} + \langle \nu', f' \rangle_{\boldsymbol{K}} . \end{aligned} \tag{4.8}$$

Then by Lemmas 3.2, 3.3 and (4.7), we have

$$\Phi^{(m+2)} = \left(\langle \nu', f' \rangle_{\boldsymbol{K}} \right)^{(m)} = \left(a\lambda + \sum_{l=1}^{n-1} b_l \langle f_{z_l}, f' \rangle_{\boldsymbol{K}} \right)^{(m)} = a\lambda^{(m)} \tag{4.9}$$

at $(\mathbf{0}, 0)$ for $m = 0, \ldots, k$, where we used (4.6) for $m = k$. Since $\lambda^{(m)}(p) = 0$ $(m \leq k-1)$ and $\lambda^{(k)}(p) \neq 0$, (4.5) yields the first part of the lemma.

Next, we show the second part: Differentiating (4.8) with respect to z_j

$$\Phi''_{z_j} = \left\langle \nu_{z_j}, f - (X, Y) \right\rangle_{\boldsymbol{K}} + \left\langle \nu'', f_{z_j} - \boldsymbol{e}_{j+2} \right\rangle_{\boldsymbol{K}} + \frac{\partial}{\partial z_j} \langle \nu', f' \rangle_{\boldsymbol{K}} ,$$

where $\boldsymbol{e}_{j+2}$ is defined in (3.2). Like as in (4.9), applying Lemma 3.2, we have

$$\Phi^{(m)}_{z_j}(\mathbf{0}, 0) = a\lambda^{(m-2)}_{z_j} = \begin{cases} 0 & (j > m-1), \\ a(p)\lambda^{(m-2)}_{z_{m-1}}(p) \neq 0 & (j = m-1). \end{cases}$$

Since $\Phi_{x_j} = \langle \nu, -\boldsymbol{e}_j \rangle_{\boldsymbol{K}} = -\nu^j$ for $j = 1, 2$, the Jacobian matrix of $(\Phi, \Phi', \ldots, \Phi^{(k)})$ at the origin is written as

$$\begin{pmatrix} -\nu^1 & -\nu^2 & 0 & 0 & \ldots & 0 & \ldots & 0 \\ -(\nu^1)' & -(\nu^2)' & 0 & 0 & \ldots & 0 & \ldots & 0 \\ * & * & a\lambda_{z_1} & 0 & \ldots & 0 & \ldots & 0 \\ * & * & * & a\lambda'_{z_2} & \ldots & 0 & \ldots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ * & * & * & * & * & a\lambda^k_{z_{k-1}} & \ldots & 0 \end{pmatrix},$$

which is of rank $k+1$. □

Proof of Theorem 2.4. It can be straightforwardly checked that the singular point given in (1.1) is k-nondegenerate, but not $(k+1)$-singular. We now prove the converse. By Zakalyukin's theorem [11] (see also the appendix of [7]), $\mathrm{Im}(f)$ is locally diffeomorphic to the standard A_{k+1}-singularity if and only if it is $\boldsymbol{K}$-right-left equivalent to (1.1), whenever the regular points of f are dense. Thus, it is sufficient to show the versality of Φ (which implies that D_Φ is locally diffeomorphic to the standard A_{k+1}-singularity, see [4]). In fact, this is evident by Lemma 4.3. □

Remark. In [7], versal unfoldings of a cuspidal edge and a swallowtail in $\boldsymbol{R}^3$ are given. The above versal unfolding Φ is much simpler than those in [7].

Now, we give proofs of Corollaries 2.5, 2.6 and 2.8 using Theorem 2.4.

Proof of Corollary 2.5. The necessary part is obvious by Lemma 3.2. We assume that $\lambda = \lambda' = \cdots = \lambda^{(k)} = 0$, $\lambda^{(k+1)} \neq 0$ at p and the map $(\lambda, \lambda', \ldots, \lambda^{(k)})$ is non-singular at p. Then we can see that p is 1-nondegenerate, so we can take the 1-st adopted coordinate system. By Lemma 3.2, p is 2-singular if $k \geq 2$. Since (λ, λ') is full rank, p is 2-nondegenerate. We can continue this argument k times. Then we see that p is k-nondegenerate but not $(k+1)$-singular. By Theorem 2.4, we have the conclusion. □

Proof of Corollary 2.6. By Lemma 2.3, p is 2-singular if and only if $\eta \in T_pS(f) = \mathrm{Span}_{\boldsymbol{K}}\{v_1, \ldots, v_{n-1}\}$, which is equivalent to $\mu(p) = 0$. Hence p is A_2-singular point if and only if $\mu(p) \neq 0$.

Since the condition does not depend on the choice of coordinate system and a frame $\{v_1, \ldots, v_{n-1}\}$ on $S(f)$, we take a coordinate system $(x_1, \ldots, x_n)$ of U such that $x_n = \lambda$ and set $v_j = \partial/\partial x_j$ $(j = 1, \ldots, n-1)$. Then the singular set $S(f)$ is given as a hyperplane $x_n = 0$, and $\mu = \eta_n$ holds on $S(f)$, where $\eta = (\eta_1, \ldots, \eta_n)$. On the other hand, $\lambda' = d\lambda(\eta) = dx_n(\eta) = \eta_n$, and then we have $\lambda' = \mu$ on $S(f)$. Then, if p is 2-singular, it is 2-nondegenerate if and only if $d\lambda'(T_pS(f)) = d\mu(T_pS(f)) \neq 0$. Moreover, since

$$S_2 = \{q \in S(f)\,;\, \lambda'(q) = 0\} = \{q \in S(f)\,;\, \mu(q) = 0\},$$

p is 3-singular if and only if $d\mu(\eta_p) = 0$. Thus, we have a criterion for A_3-singularities. □

Proof of Corollary 2.8. Since p is 2-nondegenerate, $d\mu_p \neq 0$ holds by Corollary 2.6. Hence, in addition to the previous proof, one can take a coordinate system $(x_1, \ldots, x_n)$ on U such that $\lambda = x_n$ and $\mu = x_{n-1}$ on $S(f)$. Then the set S_2 is a linear subspace $\{(x_1, \ldots, x_{n-2}, 0, 0)\}$ near p. As seen in the previous proof, $\lambda' = \mu$ holds on $S(f)$, and hence the condition (2.5) in Corollary 2.5 is equivalent to (2.7) under the assumption that p is 2-singular. Moreover, since $\lambda'' = \mu'$ on S_2, the Jacobian matrix of the map Λ in Corollary 2.5 is computed as

$$\begin{pmatrix} \lambda_{x_1} & \cdots & \lambda_{x_{n-2}} & \lambda_{x_{n-1}} & \lambda_{x_n} \\ \lambda'_{x_1} & \cdots & \lambda'_{x_{n-2}} & \lambda'_{x_{n-1}} & \lambda'_{x_n} \\ \lambda''_{x_1} & \cdots & \lambda''_{x_{n-2}} & \lambda''_{x_{n-1}} & \lambda''_{x_n} \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ \lambda^{(k-1)}_{x_1} & \cdots & \lambda^{(k-1)}_{x_{n-2}} & \lambda^{(k-1)}_{x_{n-2}} & \lambda^{(k-1)}_{x_n} \end{pmatrix} = \begin{pmatrix} 0 & \cdots & 0 & 0 & 1 \\ 0 & \cdots & 0 & 1 & * \\ \mu'_{x_1} & \cdots & \mu'_{x_{n-2}} & * & * \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ \mu^{(k-1)}_{x_1} & \cdots & \mu^{(k-1)}_{x_{n-2}} & * & * \end{pmatrix}$$

holds at p. Hence the conclusion follows. □

Example 4.4 (Tangent developables in $\boldsymbol{K}^4$). Let $\gamma(z)$ be a $\boldsymbol{K}$-differentiable curve in $\boldsymbol{K}^4$ such that $\gamma', \gamma'', \gamma''', \gamma''''$ are linearly independent. We consider a hypersurface $f(z, u, v) = \gamma(z) + u\gamma'(z) + v\gamma''(z)$. If we set $\nu = \gamma' \wedge \gamma'' \wedge \gamma'''$, then f is a front with $\boldsymbol{K}$-normal vector ν. Furthermore, we have

$$S_1 = \{v = 0\}, \quad S_2 = \{u = v = 0\}, \quad \eta = (-1, 1, u) \quad \text{and} \quad \lambda = v.$$

By Corollary 2.5, f at $(z, u, 0)$ is $\boldsymbol{K}$-right-left equivalent to (cusp) $\times \boldsymbol{K}^2$ if and only if $u \neq 0$, and f at $(z, 0, 0)$ is $\boldsymbol{K}$-right-left equivalent to (swallowtail) $\times \boldsymbol{K}$.

5. Zigzag numbers on A-fronts

Let M^n (resp. N^{n+1}) be a manifold of dimension n (resp. $n+1$), and $f\colon M^n \to N^{n+1}$ a front. If all singularities of f are A_k-singular points ($k \leq n+1$), it is called an *A-front*. In this section, we set $\boldsymbol{K} = \boldsymbol{R}$, and discuss a topological invariant of A-fronts. Now, we assume that M^n is orientable.

For each A_k $(k \leq n+1)$ singular point on an A-front, the image of it coincides with the discriminant set $\{F = F_t = 0\}$ of (1.2). Then we can define the *inward normal vector* ν_{in} which points in the direction that the number of roots of polynomial $F(t) = 0$ increasing. For example, on the A_2-singularity $(t, x_2, \ldots, x_n) \mapsto (2t^3, -3t^2, x_2, \ldots, x_n)$, ν_{in} is defined by

$$\nu_{\mathrm{in}} = \begin{cases} -(1, t, 0, \ldots, 0) & (t > 0), \\ (1, t, 0, \ldots, 0) & (t < 0), \end{cases}$$

see Figure 5.1, left. Then, the inward direction is uniquely defined on a neighborhood of each connected component of $S(f)$. On the other hand, a front $f : M^n \to$

N^{n+1} is called a *Morin front* if there exists a C^∞-map $\hat{f}$ from an $(n+1)$-manifold into N^{n+1} such that

- $\hat{f}$ admits only Morin singularities,
- f is the restriction of $\hat{f}$ to the singular set $S(\hat{f})$.

Morin fronts are all A-fronts (see Corollary 2.11 and the appendix). Conversely, any A-fronts are locally Morin fronts. The inward direction of the Morin front f points in the direction of the image of $\hat{f}$, and we get the following:

Proposition 5.1. *A Morin front admits has a unique unit normal vector on the set of regular ponits which points in the inward direction.*

We fix a regular point $o \in M^n$ of f. A continuous map $\gamma : [0,1] \longrightarrow M^n$ is called a *loop* at o if it satisfies $\gamma(0) = \gamma(1) = o$. We fix a Riemannian metric g of N^{n+1}. A loop γ is called *co-orientable* if there exists a unit normal vector field $\nu(t) \in T_{\hat{\gamma}(t)}N^{n+1}$ of f along γ such that $\nu(0) = \nu(1)$, where $\hat{\gamma}(t) := f \circ \gamma(t)$. If γ is not co-orientable, it is called *non-co-orientable*. These two properties of loops depend only on their homotopy classes. We set

$$\rho_f(\gamma) := \begin{cases} 0 & (\text{if } \gamma \text{ is co-orientable}), \\ 1 & (\text{if } \gamma \text{ is non-co-orientable}). \end{cases}$$

Then it induces a representation $\rho_f : \pi_1(M^n) \longrightarrow \boldsymbol{Z}_2 = \boldsymbol{Z}/2\boldsymbol{Z}$. Next, we set

$$\hat{\pi}_1(M^n) := \ker(\rho_f)\big(\subset \pi_1(M^n)\big)$$

which consists of homotopy classes of co-orientable loops. We shall now construct the secondary homotopical representation $\hat{\rho}_f : \hat{\pi}_1(M^n) \longrightarrow \boldsymbol{Z}$ called the *Maslov representation* as follows: Let $f : M^n \to N^{n+1}$ be a front and

$$S(f) = \bigcup_{\alpha \in \Lambda} \Gamma_\alpha$$

be the decomposition of the singular set $S(f)$ by its connected components. Then there exists a disjoint family of open domains O_α in M^n such that $\Gamma_\alpha \subset O_\alpha$. Let $\gamma : [0,1] \to M^n$ be a co-orientable loop at $o \notin S(f)$. We set

$$S_\gamma := \{t \in [0,1]\,;\, \gamma(t) \in S(f)\}.$$

For each $t \in S_\gamma$, there exists an open interval $I_t (\subset (0,1))$ such that $\gamma(I_t) \subset O_\alpha$ for some $\alpha \in \Lambda$. Since S_γ is compact, there exists a finite number of open intervals $I_1 = (t_1^-, t_1^+), \ldots, I_m = (t_m^-, t_m^+)$ such that $S_\gamma \subset I_1 \cup \cdots \cup I_m$. If $I_k \cap I_{k+1}$ is non-empty, we can replace I_k and I_{k+1} by $I_k \cup I_{k+1}$. After these operations, each I_k is still contained in some O_α, since $\{O_\alpha\}_{\alpha \in \Lambda}$ is disjoint. Then we may assume that

$$0 < t_1^- < t_1^+ < t_2^- < \cdots < t_m^- < t_m^+ < 1.$$

We fix a unit normal vector $\nu_0 \in T_{\hat{\gamma}(o)}N^{n+1}$ at o. (If f admits a globally defined smooth unit normal vector field ν on M^n, one of the canonical choices is $\nu_0 = \nu(0)$.) Let ν_j $(j = 1, 2, \ldots, m)$ be the inward unit normal vector at $t = t_j^+$. Then we can take a continuous unit vector field $\hat{\nu}_j(t)$ $(t_j^+ \le t \le t_{j+1}^-)$ along $\gamma|_{[t_j^+, t_{j+1}^-]}$ such that


$$\hat{\nu}_i(t_j^+) = \nu_j \qquad (i = 0, 1, 2, \ldots, m),$$

where $t_0^+ = 0$ and $t_{m+1}^- = 1$. We set $\varepsilon_0 = +1$ and

$$\varepsilon_j := \begin{cases} \varepsilon_{j-1} & (\text{if } \hat{\nu}_{j-1}(t_j^-) \text{ is inward}) \\ -\varepsilon_{j-1} & (\text{if } -\hat{\nu}_{j-1}(t_j^-) \text{ is inward}) \end{cases} \qquad (j = 1, 2, \ldots, m+1),$$

where $\nu_{m+1} = \nu_0$. Then we get a sequence

$$\varepsilon_0,\ \varepsilon_1,\ \varepsilon_2,\ \ldots\ ,\ \varepsilon_m,\ \varepsilon_{m+1}. \tag{5.1}$$



The inward normal Zigzag number

FIGURE 5.1. Zigzag number of a plane curve

Since M^n is orientable, m is an odd integer. By the co-orientability of γ, $\varepsilon_0 = \varepsilon_{m+1} = +1$ holds. For simplicity, we write $\varepsilon_j = +$ (resp. $-$) if $\varepsilon_j = +1$ (resp. -1). If there are two successive same signs $++$ or $--$, we cancel them. Repeating this cancellation, we get a sequence which has no successive signs

$$+ - \cdots + - + \,.$$

We shall call this reduction the *normalization* of the sequence (5.1). Then the number of $-$ after the normalization is called the *zig-zag number* of the co-orientable loop γ. (The definition of zig-zag number for surfaces in $\boldsymbol{R}^3$ is given in [8]. See also [10].) For example, we set $M = S^1$ and $N = \boldsymbol{R}^2$ and consider a front as in Figure 5.1, right. Then this closed curve induces a sequence

$$+ - + - + - - + - - ++$$

which can be reduced to

$$+ - +(-(+(--)+)-) - (++) \quad \text{that is} \quad + - + -$$

and thus the zig-zag number in this case is 2. We denote by $z(\gamma)$ the zig-zag number of the co-orientable loop γ. (Global properties and references of zig-zag numbers of planar curves are given in [1].)

Proposition 5.2. *The zig-zag number $z(\gamma)$ depends only on the homotopy class of γ, and is stable under the deformation of an A-front.*

Proof. Suppose $[\gamma_0] = [\gamma_1]$ in $\pi_1(M^n)$. Then there exists a homotopy $\Gamma : [0,1] \times [0,1] \to M^n$ such that $\Gamma(t,0) = \gamma_0(t)$ and $\Gamma(t,1) = \gamma_1(t)$. We set

$$S_\Gamma := \{(t,s) \in [0,1] \times [0,1] \,;\, \Gamma(t,s) \in S(f)\}.$$

For each $(t,s) \in S_\Gamma$, there exists an open neighborhood $W_{t,s} \subset [0,1] \times [0,1]$ of (t,s) such that $\Gamma(W_{t,s}) \subset O_\lambda$ for some $\lambda \in \Lambda$. Since $[0,1] \times [0,1]$ is compact, it is covered by just finitely many such open subsets. Then the standard homotopy argument yields the homotopy invariance $z(\gamma_0) = z(\gamma_1)$. □

A closed C^1-regular curve $\gamma : [0,1] \to M^n$ starting at o is called a *null loop* if $\gamma'(t)$ $(t \in [0,1])$ belongs to the kernel of df whenever $\gamma(t) \in S(f)$. One can easily show that, for each homotopy class of the fundamental group $\pi_1(M^n)$, there exists a null C^1-loop which represents the homotopy class. Moreover, we may assume that $\gamma(t)$ passes through only finitely many A_2-singular points. Moreover, we may assume that $\gamma(t)$ passes through only finitely many A_2-singular points.

Suppose that $\gamma(t)$ is a co-orientable null loop. Then there exists a continuous unit normal vector field $\nu(t)$ along $\gamma(t)$ such that $\nu(0) = \nu_0$. Since γ is co-orientable,

$\nu(1) = \nu_0$ holds. As in [10], we define a continuous map of $[0,1] \setminus S_\gamma$ into $P^1(\boldsymbol{R})$ by

$$C_\gamma : t \longmapsto [g(\hat{\gamma}'(t), \hat{\gamma}'(t)) : g(\nu'_\gamma(t), \hat{\gamma}'(t))],$$

which can be extended continuously across S_γ. We call this map $C_\gamma : [0,1] \to P^1(\boldsymbol{R})$ *the normal curvature map*. In fact, when $\gamma(t)$ is a regular point of f, $-g(\nu'_\gamma, \hat{\gamma}')/g(\hat{\gamma}', \hat{\gamma}')$ is exactly the normal curvature of f along γ. This is independent of the orientation of γ, but its sign depends on the initial choice of the unit normal vector $\pm\nu_0$ at o. We denote the rotation index of the map C_γ by $\mu_f(\gamma)$. The absolute value of $\mu_f(\gamma)$ is called the *Maslov index* of the co-orientable null loop γ. Then by the same argument as in [10], we get the following

Proposition 5.3. *The Maslov index $|\mu_f(\gamma)|$ coincides with the zig-zag number $z(\gamma)$. In particular, we get a (Maslov) representation*

$$\hat{\rho}_f : \hat{\pi}_1(M^n) \ni [\gamma] \longmapsto \mu_f(\gamma) \in \boldsymbol{Z}.$$

Since the inward direction is defined globally for a Morin front, we have the following:

Corollary 5.4. *The Maslov representations for Morin fronts are trivial.*

Remark 5.5. Suppose that M^n and N^{n+1} are both oriented and f is co-orientable. Then, we can take a globally defined unit normal vector field ν along f which gives a direct interpretation of the sign ε_j: We denote by O^+_α the component of $O_\alpha \setminus \Gamma_\alpha$ where ν is compatible with the orientation of M^n. Then each component Γ_α of $S(f)$ is called *zig* if ν coincides with ν_{in} on O^+_α. Some global properties of zig-zag numbers are given in [8].

Appendix A. Criteria of Morin type singularities

The *A_k-Morin singularities* are map germs which are $\boldsymbol{K}$-right-left equivalent to

$$f(z_1, \ldots, z_n) = (z_1 z_n + \cdots + z_{k-1} z_n^{k-1} + z_n^{k+1}, z_1, \ldots, z_{n-1}) \qquad (k \leq n) \tag{A.1}$$

at $\mathbf{0}$ ([9], see also [5]). The A_0-Morin singularity means a regular point. The following assertion holds:

Theorem A.1. *Assume that $k \leq n$. Let U be a domain in $\boldsymbol{K}^n$, and $f\colon U \to \boldsymbol{K}^n$ a $\boldsymbol{K}$-differentiable map and $p \in S(f)$. We assume that p is a corank one singularity. Then f at p is $\boldsymbol{K}$-right-left equivalent to A_k-Morin singularity if and only if*

$$\lambda = \lambda' = \cdots = \lambda^{(k-1)} = 0,\ \lambda^{(k)} \neq 0 \text{ at } p \quad \text{and} \tag{A.2}$$

$$\Lambda := (\lambda, \lambda', \ldots, \lambda^{(k-1)}) : U \to \boldsymbol{K}^k \quad \text{is non-singular at } p. \tag{A.3}$$

Here, $\lambda = \det(f_{x_1}, \ldots, f_{x_n})$, $(x_1, \ldots, x_n)$ is the canonical coordinate system on U, $\lambda' = \tilde{\eta}\lambda (= \lambda^{(1)})$, $\lambda^{(i)} = \tilde{\eta}\lambda^{(i-1)}$ and $\tilde{\eta}$ is the extended null vector field of f.

Proof. Since the conditions (A.2) and (A.3) do not change when changing λ to $\psi\lambda$, where $\psi(p) \neq 0$, the conditions (A.2) and (A.3) are independent on the choice of local coordinate systems on the source and target. Suppose f has the form

$$f(z_1, \ldots, z_n) = \big(\mu(z_1, \ldots, z_n), z_1, \ldots, z_{n-1}\big), \tag{A.4}$$

such that

- $\mu' = \cdots = \mu^{(k)} = 0$, $\mu^{(k+1)} \neq 0$ at p ($' = \partial/\partial z_n$), and
- the map germ of $(\mu', \ldots, \mu^{(k)})$ is nonsingular at p.

Then f is an A_k-Morin singularity (see [5, page 177]). Though the proof of this fact in [5] is given for $\boldsymbol{K} = \boldsymbol{R}$, the argument works also for the case of $\boldsymbol{K} = \boldsymbol{C}$. By (A.4), we have $\lambda = \mu'$, and $\eta = \partial_{z_n}$ holds. This implies the theorem. □

The same type of assertion as in Corollaries 2.6 and 2.8 hold for A_k-Morin singularities.

Acknowledgements. The authors thank Shyuichi Izumiya and Goo Ishikawa for fruitful discussions and valuable comments.

References

[1] V. I. Arnol'd, Topological Invariants of Plane Curves and Caustics, University Lecture Series **5**, Amer. Math. Soc. (1991).
[2] V. I. Arnol'd, S. M. Gusein-Zade and A. N. Varchenko, Singularities of differentiable maps, Vol. 1, Monographs in Math. **82**, Birkhäuser (1984).
[3] J. M. Boardman, *Singularities of differentiable maps*, Publ. Math. I.H.E.S. **33**, (1967), 21–57.
[4] J. W. Bruce and P. J. Giblin, *Curves and singularities*, Camb. Univ. Press (1992).
[5] M. Golubitsky and V. Guillemin, *Stable mappings and their singularities*, Graduate Texts in Math. **14**, Springer-verlag, 1973.
[6] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry I,* John Wiley & Sons Inc., 1963.
[7] M. Kokubu, W. Rossman, K. Saji, M. Umehara and K. Yamada, *Singularities of flat fronts in hyperbolic* 3*-space*, Pacific J. Math. **221** (2005), no. 2, 303–351.
[8] R. Langevin, G. Levitt and H. Rosenberg, *Classes d'homotopie de surfaces avec rebroussements et queues d'aronde dans* $\mathbb{R}^3$, Canad. J. Math. **47** (1995), 544–572.
[9] B. Morin, *Forms canonique des singularités d'une application différentiable*, C. R. Acad. sci. Paris **260** (1965), 5662–5665.
[10] K. Saji, M. Umehara and K. Yamada, *The geometry of fronts*, to appear in Ann. of Math., math.DG/0503236.
[11] V. M. Zakalyukin, *Reconstructions of fronts and caustics depending on a parameter and versality of mappings.* J. Sov. Math., **27** (1984), 2713–2735.

(Saji) Department of Mathematics, Faculty of Educaton, Gifu University, Yanagido 1-1, Gifu 151-1193, Japan
E-mail address: ksaji@gifu-u.ac.jp

(Umehara) Department of Mathematics, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan
E-mail address: umehara@math.sci.osaka-u.ac.jp

(Yamada) Faculty of Mathematics, Kyushu University, Higashi-ku, Fukuoka 812-8581, Japan
E-mail address: kotaro@math.kyushu-u.ac.jp